Parkinson's Tremors: What Causes Them and What Helps

Ashal Abid Satti

1st Year MBBS, Islamabad Medical and Dental College, Islamabad, Pakistan

Key points:

- Introduction
- Pathophysiology
- Complications
- Diagnosis
- Management
- Conclusion

Introduction

Parkinson's disease (PD), also known as parkinsonism, is a progressive neurodegenerative disorder that primarily affects movement. It is characterized by motor symptoms such as tremors, rigidity, and bradykinesia. Among these, tremors are often the earliest and most noticeable feature, frequently leading patients to seek medical attention. However, tremor severity, type, and response to treatment vary significantly among individuals.

Recent advances have shown that Parkinsonian tremors are not solely due to degeneration of dopaminergic neurons in the substantia nigra. Instead, they arise from dysfunction within interconnected neural circuits involving the basal ganglia, thalamus, and cerebellum. These circuits play a central role in coordinating movement, and their disruption leads to the abnormal rhythmic activity seen in tremor.

Pathophysiology

Parkinsonian tremors develop due to abnormal rhythmic activity within the motor control systems of the central nervous system. Although degeneration of dopaminergic neurons in the substantia nigra is a key feature of Parkinson's disease, it does not fully account for tremor generation. The basal ganglia thalamocortical circuit is primarily responsible for rest tremors. This tremor typically occurs at a frequency of four to six hertz and is associated with synchronized neuronal activity in structures such as the globus pallidus and thalamus.

In contrast, the cerebellothalamic circuit contributes to re-emergent tremor, which develops after a brief latency when a posture is maintained. This type of tremor often shows reduced responsiveness to dopaminergic therapy, suggesting the involvement of non dopaminergic pathways.

Other neurotransmitters, including serotonin and noradrenaline, also influence tremor amplitude and frequency. This supports the concept that Parkinsonian tremor is a network disorder involving multiple pathways rather than a single localized defect.

Clinical Features

Tremors in Parkinson's disease present in different forms, each with distinct characteristics. The most common type is rest tremor, which occurs when the affected body part is relaxed and decreases with voluntary movement. Re emergent tremors appear

after a short delay when maintaining posture and may be confused with postural tremors.

Although less common, postural, and kinetic tremors can also occur in some patients. A classic feature of Parkinsonian tremor is the pill rolling movement of the fingers, most often seen in the hands. Tremor usually begins on one side of the body and often remains asymmetric.

Patients with tremor dominant Parkinson's disease tend to have slower progression of overall motor disability. Despite this, tremor can significantly interfere with daily activities, particularly those requiring fine motor skills such as writing, buttoning clothes, and handling small objects.

Diagnosis & Supporting Investigations

The diagnosis of Parkinson's disease is primarily clinical and is based on a detailed history and neurological examination. Key features include unilateral onset of symptoms, presence of rest tremor, and associated signs such as rigidity and bradykinesia. The progression of symptoms over time further supports the diagnosis.

Although clinical assessment remains the gold standard, several supportive tools are increasingly used. DaTSCAN imaging helps differentiate Parkinson's disease from essential tremor by evaluating dopamine transporter activity.

Wearable devices and smartphone based accelerometers are also being used to objectively assess tremor characteristics. Clinical rating scales, such as the Movement Disorder Society Unified Parkinson's Disease Rating Scale, are useful for evaluating disease severity and progression. In addition, machine learning based tools are showing promise in improving early detection and diagnostic accuracy.

Management of Parkinson's Disease Associated Tremors

Management involves a combination of pharmacological therapy, surgical intervention, and emerging technologies. The choice of treatment depends on symptom severity, patient age, and response to previous therapies. A personalized approach is essential because tremor response varies among individuals.

Levodopa remains the most effective treatment and is commonly used as first line therapy. However, its effect on tremor may be inconsistent, especially when non dopaminergic pathways are involved. Additional therapies are often required to achieve better symptom control.

- **Levodopa therapy** is the mainstay of treatment, although tremor response may vary
- **Dopamine agonists** are useful in early stages and may delay levodopa use
- **Anticholinergic drugs** may benefit younger patients with prominent tremor
- **Deep brain stimulation** is effective for patients with drug resistant tremor
- **Focused ultrasound** provides a non invasive treatment alternative
- **Adaptive DBS** and **gene based therapies** are emerging treatment options

Surgical interventions such as deep brain stimulation target specific brain regions involved in tremor generation and can provide significant relief in selected patients. Non invasive approaches like focused ultrasound are increasingly used due to their favorable safety profile.

Conclusion

Tremor in Parkinson's disease is a complex symptom resulting from dysfunction of multiple interconnected neural networks. It cannot be explained by a single

pathological mechanism, which accounts for its variability in presentation and response to treatment.

Advances in understanding the underlying mechanisms, along with improvements in diagnostic tools and treatment strategies, have significantly enhanced the management of Parkinson's disease. Ongoing research is focused on developing more targeted and individualized therapies to further improve patient outcomes and quality of life.

References

1. Dirkx M, Helmich R. Pathophysiology and management of parkinsonian tremor. Seminars in Neurology. 2017 May 16;37(02):127–34. doi:10.1055/s-0037-1601558
2. Ortega-Robles E, Arias-Carrión O. Tremor: clinical frameworks, network dysfunction and therapeutics. Brain Sciences. 2025 Jul 27;15(8):799
3. 1. Deuschl G, Raethjen J, Baron R, Lindemann M, Wilms H, Krack P. The pathophysiology of parkinsonian tremor: A review. Journal of Neurology. 2000 Oct 6;247(S5). doi:10.1007/pl00007781
4. 1. Picillo M, Lozano AM, Kou N, Puppi Munhoz R, Fasano A. Programming deep brain stimulation for parkinson's disease: The Toronto Western Hospital algorithms. Brain Stimulation. 2016 May;9(3):425–37. doi:10.1016/j.brs.2016.02.004